

CME Observation during the Total Solar Eclipse on November 14, 2012 (2)

Osamu OHGOE

Amateur Astronomer -- Tokyo Japan; sol-eclipse.ohgoe@nifty.com

(2013.06.13)

ABSTRACT

During the total solar eclipse on November 14, 2012 the white-light corona was observed in the northern Australia and Pacific Ocean. I took many frames of white-light corona on the ship "Pacific Venus" cruising about 500 km north from New Zealand. In the frames I took, the CME can be discerned but in frames taken at Australia, 35 minutes before, there is no image of CME. Then, I concluded transverse velocity of the CME was 220~580km/s and the transverse acceleration was 2.5~3.7 km/s².

1. Introduction

In order to measure coronal motions, multi point observations of total solar eclipse had planed many times. For example, members of Tokyo University of Science Astronomical Club observed the total solar eclipse on February 16, 1980 in Kenya and India. The time difference between two sites is about 110 minutes. They caught movement of two loops and one small streamer of corona. The transverse velocity of each was about 20km/s, 7~8km/s and 23km/s [1](see Figure 1). Generally, the velocity of the change of the coronal structures is small. In order to catch them, we need two observation sites having some time difference.




Figure 1 Movement of structures of the corona on 1980 February 16

There is a fast-moving phenomenon in the corona like a CME (coronal mass ejection). It is not unusual to occur some CMEs during a total solar eclipse [2](see Figure 2). But it is very rare that we can see the motion of the CME at one observation site.

2. Basic Information on the 2012 November 14 eclipse observation

The path of totality stretched from northern Australia to the Pacific Ocean. In the Pacific Ocean




Figure 2 Computer-processed image of the corona taken by Osamu OHGOE, 2010 July 11 at HAO atoll, French Polynesia.

the Moon's shadow made no landfall. Our ship "Pacific Venus" left Noumea of New Caledonia on Nov. 12 for Auckland in New Zealand. When we caught the total solar eclipse on the way, the location of the ship was $173^{\circ}04'32''E \sim 173^{\circ}04'23''E$ and $30^{\circ}02'04''S \sim 30^{\circ}02'02''S$ [3].




Figure 3 The location of our ship during the total solar eclipse, GPS Data provided by Jun Nakazawa

I observed the eclipse with one telescope 100mm in diameter and 512mm in a focal length, and took 209 frames of the corona. With 33 frames among them, I got 5 images of the fine structure of the white-light corona processed by Shiota's method [4].

3. Movement of the CME

Figure 4~6 illustrates image A, C and E. Each image was taken at 21:14:11 UT, 21:15:08 UT and 21:16:12 UT. I chose 3 points in the front of the CME and Figure 7 illustrate them. The radius of the





Figure 4 Image A (top left)

Figure 5 Image C (top right)

Figure 6 Image E (bottom)

Figure 7 Choice 3 points
in the front of the CME

Sun is 6.960×10^5 km. The semi diameter of the Sun is 969 seconds of arc. The semi diameter of the Moon is 1017 seconds of arc.

The provisional values of transverse velocity and acceleration of the CME were calculated as follows.

I used three images for a calculation.

Image No.6229	21:14:13(UT)	1/250 second exposure
Image No.6277	21:15:08(UT)	1/250 second exposure
Image No.6329	21:16:13(UT)	1/250 second exposure

Point 1	transverse velocity	transverse acceleration
No.6229	$(3.7 \pm 0.3) \times 10^2$ km/s	
No.6277		3.2 ± 0.8 km/s ²
No.6329	$(5.7 \pm 0.2) \times 10^2$ km/s	
Point 2	transverse velocity	transverse acceleration
No.6229	$(2.2 \pm 0.3) \times 10^2$ km/s	
No.6277		2.5 ± 0.8 km/s ²
No.6329	$(3.7 \pm 0.2) \times 10^2$ km/s	
Point 3	transverse velocity	transverse acceleration
No.6229	$(3.5 \pm 0.3) \times 10^2$ km/s	
No.6277		3.7 ± 0.8 km/s ²
No.6329	$(5.8 \pm 0.2) \times 10^2$ km/s	

These speed is the ingredient which is perpendicular to the line of sight. This value seems to be a proper value for the speed of CME even if I take the factor into account.

I think that the sharpening of the image and improvement of the time resolution are important by the future observation.

REFERENCES

- [1] Ohgoe, O. ,et al. 1980, Proceedings of The annual meeting of Astronomical Society of Japan in 1980 autumn (Japanese)
- [2] Ohgoe, O. ,et al. 2010, Reports of the Total Solar Eclipse 2010 July 11 at French Polynesia ~ DVD (Japanese)
- [3] Ohgoe, O. ,et al. 2013, Reports of the Total Solar Eclipse 2012 November 14 ~ DVD (Japanese)
- [4] Ohgoe, O. , Shiota, K. , 2012, Chasing Shadows: An Observer's Guide to Solar Eclipses (Japanese), Seibundo Shin Kousha